arXiv:1603.03526v1 [math.CO] 11 Mar 2016

Cycles in graphs of fixed girth with large size

József Solymosi*, Ching Wong†

Abstract

Consider a family of graphs having a fixed girth and a large size. We give an optimal lower asymptotic bound on the number of even cycles of any constant length, as the order of the graphs tends to infinity.

1 Introduction

All graphs we consider in this article are simple graphs. We denote the size of G by $e(G)$ and the order of G by $v(G)$. A j-path in G is a path of length j in G. A j-cycle in G is a cycle of length j in G, and it is called an even cycle if j is even. The *girth* of a graph G is the length of the shortest cycles in G. For $x \in V(G)$, let $\Gamma_G^k(x) = \Gamma^k(x)$ denote the set of vertices of G having distance exactly k from the vertex x.

In the following, the big-O notations $f(n) = O(g(n))$ are understood as $f(n) = O(g(n))$ as $n \to \infty$, where n denotes the order of a graph. The same applies to Θ.

It is easy to see that a graph having large girth cannot have too many edges. The famous Erdős girth conjecture asserts the existence of graphs of any given girth with a size of maximum possible order.

Conjecture 1 (Erdős girth conjecture)**.** *For any positive integer m, there exist a constant $c > 0$ depending only on m and a family of graphs $\{G_n\}$ such that $v(G_n) = n$, $e(G_n) \geq cn^{1+1/m}$ and girth$(G_n) > 2m$.*

*Research was supported by NSERC, ERC-AdG. 321104, and OTKA NK 104183 grants.

†Research was supported by FYF (UBC).

Indeed, such size is maximum by the result of Bondy and Simonovits [2], in which an explicit constant is given. They showed that a graph G_n of order n with $\text{girth}(G_n) > 2m$ has a size less than $100mn^{1+1/m}$.

This conjecture has been proved true for $m = 1, 2, 3, 5$. See [5], [3], [1] and [7]. For a general m, Sudakov and Verstraëte [6] showed that if such graphs exist, then they contain at least one cycle of any even length between and including $2m + 2$ and Cn, for some constant $C > 0$.

For $\ell > m$, by counting the number of $(2\ell - 2m)$-paths, one can show that the number of 2ℓ-cycles in such graphs has an order not greater than $O(n^{2\ell/m})$, provided that $\deg_{G_n}(x) = \Theta(n^{1/m})$ for any vertex x in G_n. We will see in Section 2 that in the asymptotic case, one can assume this without loss of generality. This suggests the definition of *almost regularity* given in Section 2.

In this article, we give a lower bound on the number of 2ℓ-cycles when $\ell = O(1)$, and conclude that the number of 2ℓ-cycles is $\Theta(n^{2\ell/m})$. The precise statement is the following.

Theorem 2. *For any real number $c > 0$ and integers M, m with $M > m \geq 2$, there exist a constant $\alpha > 0$ and an integer N such that if $\{G_n\}$ is a family of graphs satisfying $v(G_n) = n$, $e(G_n) \geq cn^{1+1/m}$ and girth$(G_n) > 2m$, then for $n \geq N$ and $m + 1 \leq \ell \leq M$, the number of 2ℓ-cycles in G_n is at least $\alpha n^{2\ell/m}$.*

We proceed as follows. In Section 2, we show that by adjusting the threshold N in our theorem, we can further assume that the graphs have some nice properties, namely bipartite and almost regular. In Section 3, we count the number of short even cycles in G_n up to length $4m$. Finally, in Section 4, we entend the argument to longer cycles, completing the proof of the main theorem.

2 Reduction to a simpler case

In this section, we show that it suffices to consider only bipartite graphs which are almost regular, defined as follows.

Definition 3. Suppose $\{G_n\}$ is a family of graphs with $v(G_n) = n$ and $\text{girth}(G_n) > 2m$, we say that $\{G_n\}$ is *almost regular* if there exist $c_1, c_2 > 0$ such that $c_1 n^{1/m} \leq \deg(x) \leq c_2 n^{1/m}$ for any vertex $x \in V(G_n)$.

It is a well-known fact that any graph has a bipartite subgraph with at least half of its edges. It remains to construct subgraphs whose maximum and minimum degree is of order $n^{1/m}$. To achieve this, we repeatedly apply a theorem of Bondy and Simonovits [2], which states that if an n-vertex graph G_n has girth larger than $2m$, then G_n has less than $100mn^{1+1/m}$ edges.

First we delete vertices of small degree.

Lemma 4. *For any real number $c > 0$ and integer $m \geq 2$, there exists a constant $\beta > 0$ such that any bipartite graph G of order n, size at least $cn^{1+1/m}$ and girth larger than $2m$ has a subgraph H having at least βn vertices, at least $\frac{9c}{10}n^{1+1/m}$ edges and minimum degree at least $\frac{c}{10}n^{1/m}$.*

Proof. Let $G = H_0$. For $i \geq 1$, inductively define H_i to be the subgraph of H_{i-1} induced by all vertices having degree at least $\frac{c}{10}n^{1/m}$ in H_{i-1}. Then for all i we have $e(H_i) \geq e(H_{i+1})$ and

$$\begin{aligned} e(H_i) &\geq e(H_0) - (v(H_0) - v(H_i))\frac{c}{10}n^{1/m} \\ &\geq cn^{1+1/m} - n\frac{c}{10}n^{1/m} \\ &= \frac{9c}{10}n^{1+1/m}, \end{aligned}$$

and so there exists some $j \geq 0$ such that $H_i = H_j$ for all $i \geq j$.

Set $H = H_j$, which has girth larger than $2m$ and size at least $\frac{9c}{10}n^{1+1/m}$. Then,

$$\frac{9c}{10}n^{1+1/m} < 100m \cdot v(H)^{1+1/m},$$

or $v(H) \geq \beta n$, where

$$\beta = \left(\frac{9c}{1000m}\right)^{m/(m+1)}.$$

□

Lemma 5. *For every real number $c > 0$ and integer $m \geq 2$, there exists a constant $\gamma > 0$ such that any bipartite graph G of order n, size at least $cn^{1+1/m}$ and girth larger than $2m$ has a subgraph H having at least $\frac{n}{2}$ vertices, at least $\frac{c}{4}n^{1+1/m}$ edges and maximum degree at most $\gamma n^{1/m}$.*

Proof. For any $\gamma > 0$, let S_γ be the set of vertices of G having degree at least $\gamma n^{1/m}$, and let T_γ be the remaining vertices of G. We want to find a γ so

large that H can be chosen as the subgraph $G[T_\gamma]$ of G induced by T_γ. It suffices to find γ large enough so that $e(G[S_\gamma]) < e(G)/4$, $e(T_\gamma, S_\gamma) < e(G)/2$ and $v(G[T_\gamma]) \geq n/2$.

Since both G and $G[S_\gamma]$ have girth larger than $2m$, we can apply the result of [2] twice to obtain

$$\begin{aligned} e(G[S_\gamma]) &< 100m|S_\gamma|^{1+1/m} \\ &\leq 100m\left(\frac{2e(G)}{\gamma n^{1/m}}\right)^{1+1/m} \\ &\leq 100m\left(\frac{2\cdot 100mn^{1+1/m}}{\gamma n^{1/m}}\right)^{1+1/m} \\ &= \left(\frac{2}{\gamma}\right)^{1+1/m}(100m)^{2+1/m}n^{1+1/m}. \end{aligned}$$

To satisfy the first condition, we choose γ large enough so that $e(G[S_\gamma]) < \frac{e(G)}{4} < \frac{100m}{4}n^{1+1/m}$, or

$$\gamma > 2\cdot 100m\cdot 4^{m/(m+1)} > 400m. \tag{1}$$

The second condition can be obtained via its contrapositive. Suppose $e(S_\gamma, T_\gamma) \geq \frac{c}{2}n^{1+1/m}$. Let G_γ be the subgraph of G induced by $E(S_\gamma, T_\gamma)$. Then apply Lemma 4 to G_γ, we get a subgraph H_γ of G_γ having $\nu \geq \left(\frac{9c}{2000m}\right)^{m/(m+1)} n$ vertices and minimum degree at least $\frac{c}{20}n^{1/m}$. Now, we consider the subgraph G'_γ of G deleting the edges in $E(T_\gamma)$. Then, in G'_γ, every vertex in $S_\gamma \cap V(H_\gamma)$ still has degree at least $\gamma n^{1/m}$ and every vertex in $T_\gamma \cap V(H_\gamma)$ has degree at least $\frac{c}{20}n^{1/m}$. Note that any m-path in G'_γ has at most $\lfloor m/2 \rfloor$ internal vertices in T_γ, therefore the number of m-paths in G'_γ is at least

$$\frac{1}{2}\nu\left(\frac{c}{20}n^{1/m}\right)^{\lfloor m/2\rfloor}\left(\gamma n^{1/m}\right)^{\lceil m/2\rceil} \geq \frac{1}{2}\left(\frac{9c}{2000m}\right)^{m/(m+1)}\left(\frac{c\gamma}{20}\right)^{\lfloor m/2\rfloor} n^2.$$

But the number of m-paths in G cannot be larger than n^2, since otherwise there is a pair of vertices being the endpoints of two m-paths, contradicting the girth of G is larger than $2m$. Hence,

$$\frac{1}{2}\left(\frac{9c}{2000}\right)^{m/(m+1)}\left(\frac{c\gamma}{20}\right)^{\lfloor m/2\rfloor} < 1,$$

or

$$\gamma < \frac{20}{c}\left(2\left(\frac{2000}{9c}\right)^{m/(m+1)}\right)^{1/\lfloor m/2\rfloor} < \frac{40}{c}\left(\frac{2000}{9c}\right)^{3/(m+1)}.$$

Therefore, if

$$\gamma > \frac{40}{c}\left(\frac{2000}{9c}\right)^{3/(m+1)},$$

then $e(S_\gamma, T_\gamma) < \frac{c}{2}n^{1+1/m} \leq \frac{e(G)}{2}$.

Finally, since $|S_\gamma| \leq \frac{200m}{\gamma}n$, the third condition is fulfilled by (1). This finishes the proof. □

3 Counting short cycles

From now on, we suppose that G is a bipartite n-vertex graph having at least $cn^{1+1/m}$ edges with girth larger than $2m$, such that for some constants $c_1, c_2 > 0$, there holds $c_1 n^{1/m} \leq \deg_G(x) \leq c_2 n^{1/m}$ for any vertex $x \in V(G)$.

In this section, we give a lower bound on the number of the 2ℓ-cycles in G, for each $m+1 \leq \ell \leq 2m$.

We first sketch the idea. Let x be a vertex in G. Suppose we have a path of odd length k in $\Gamma_G^m(x) \cup \Gamma_G^{m+1}(x)$ with endpoints $w_0 \in \Gamma_G^m(x)$ and $w_k \in \Gamma_G^{m+1}(x)$. For each neighbor y in $\Gamma_G^m(x)$ of w_k, the four paths joining x to w_0, w_0 to w_k, w_k to y, and y to x form a closed walk, which contains a cycle, as shown in Figure 1. We show in Section 3.1 that generically these paths are internally disjoint, i.e. the length of the cycle is $2m+k+1$. Then we count in Section 3.2 the number of such paths and the number of neighbours of w_k. Finally, we obtain the desired lower bound in Section 3.3.

3.1 Internally disjoint closed walk

Note that for distinct $i, j \leq m+1$, we know that $\Gamma_G^i(x) \cap \Gamma_G^j(x)$ is empty because G is bipartite and has girth larger than $2m$. In particular, the subgraph G_x of G induced by the vertices $\Gamma_G^m(x) \cup \Gamma_G^{m+1}(x)$ is bipartite with bipartition $\{\Gamma_G^m(x), \Gamma_G^{m+1}(x)\}$. Hence, any path of odd length in G_x has one endpoint in $\Gamma_G^m(x)$ and the other endpoint in $\Gamma_G^{m+1}(x)$.

For $1 \leq i \leq m$ and any vertex $w \in \Gamma_G^i(x)$, there is a unique (x, w)-path P_w^x of length i in G. Note that for any two vertices $y_1, y_2 \in \Gamma_G^m(x)$, the intersection of the paths $P_{y_1}^x$ and $P_{y_2}^x$ must be a path, of which x is an



Figure 1: The four paths form a closed walk, which contains a cycle

endpoint. The following lemma guarantees that if $w_0 \in \Gamma_G^m(x)$ and $w_k \in \Gamma_G^{m+1}(x)$, there is at most one neighbour $u \in \Gamma_{G_x}^1(w_k)$ so that $P_{w_0}^x$ and P_u^x intersect internally, see Figure 2.



Figure 2: Other neighbours of w_k give internally disjoint paths

Lemma 6. *Suppose two vertices $y_1, y_2 \in \Gamma_G^m(x)$ share a common neighbour $w \in \Gamma_G^{m+1}(x)$, then the paths P_{y_1} and P_{y_2} are internally disjoint.*

Proof. Suppose the paths $P_{y_1}^x$ and $P_{y_2}^x$ intersects internally, then their intersection must be a path of length $L \geq 1$, with endpoints x and v, for some



Figure 3: A cycle of length at most $2m$ formed

$v \in \Gamma_G^L(x)$. Thus, the union of the paths $P_{y_1} \backslash P_v$, $P_{y_2} \backslash P_v$ and the edges (y_2, w), (w, y_1) is a cycle of length $2(m - L) + 2 \leq 2m$ in G, as in Figure 3, contradiction. □

Now, given a path $P = (w_0, w_1, \ldots, w_k)$ of odd length $k \leq 2m - 1$ in G_x with $w_0 \in \Gamma_G^m(x)$ and $w_k \in \Gamma_G^{m+1}(x)$. Note that $V(P) \cap \Gamma_{G_x}^1(w_k) = \{w_{k-1}\}$ as girth$(G_n) > 2m$. Let $y \in \Gamma_{G_x}^1(w_k)$. As shown in Figure 4, the four paths $P_{w_0}^x$, P, (w_k, y) and P_y^x contain a cycle of length $2m + k + 1$, with at most two exceptions, namely $y = w_{k-1}$ and $y = u$.



Figure 4: Most neighbours of w_k give cycles of length $2m + k + 1$

3.2 Number of paths in G_x

Note that in G_x, the minimum degree can be as small as 1. Instead of counting the number of paths of a given length in G_x, we work with a subgraph of G_x having large minimum degree. We adopt the result from Section 2.

It is easy to see that $v(G_x) \le n$ and

$$e(G_x) \ge \left|\Gamma_G^m(x)\right| c_1 n^{1/m} \ge \left(c_1 n^{1/m}\right)^{m+1} = c_1^{m+1} n^{1+1/m}.$$

Using Lemma 4, we obtain a bipartite subgraph H_x of G_x having order at least $\left(\frac{9c_1^{m+1}}{1000m}\right)^{1/(1+m)} n$, size at least $\frac{9c_1^{m+1}}{10} n^{1+1/m}$, and $\frac{c_1^{m+1}}{10} n^{1/m} \le \deg_{H_x}(u) \le c_2 n^{1/m}$, for any vertex $u \in V(H_x)$, with bipartition $\{A_x, B_x\}$, where $A_x = \Gamma_G^m(x) \cap H_x$, and $B_x = \Gamma_G^{m+1}(x) \cap H_x$.

Lemma 7. *Let k be an odd number satisfying $1 \le k \le 2m-1$. The number of k-paths in G_x is at least*

$$\frac{9c_1^{(m+1)(k+1)}}{10^{k+1} c_2} n^{1+k/m}.$$

Proof. The result follows from

$$|B_x| \ge \frac{e(H_x)}{c_2 n^{1/m}} \ge \frac{9c_1^{m+1}}{10c_2} n$$

and that the number of k-paths in G_x is at least

$$|B_x| \left(\frac{c_1^{m+1}}{10} n^{1/m}\right)^k.$$

□

3.3 Lower bound on the number of short cycles

The work in the preceding sections allows us to find a lot cycles in G. It is clear that a 2ℓ-cycle can be counted by at most 2ℓ times as each vertex of the cycle can play the role of x once.

We are ready to give a lower bound on the number of short even cycles, up to length $4m$ in G.

Proposition 8. *Let m be a positive integer. Let G be a bipartite n-vertex graph having girth larger than $2m$ and $c_1 n^{1/m} \le \deg_G(v) \le c_2 n^{1/m}$ for any vertex $v \in V(G)$, for some $c_1, c_2 > 0$. Then for $m+1 \le \ell \le 2m$, the number of 2ℓ-cycles in G is at least $\alpha_\ell n^{2\ell/m}$, where*

$$\alpha_\ell = \frac{9}{2\ell c_2}\left(\frac{c_1^{m+1}}{10}\right)^{2\ell-2m+1} > 0.$$

Proof. Using Lemma 7 with $k = 2\ell - 2m - 1$ and the observation above, the number of 2ℓ-cycles in G is at least

$$\begin{aligned}
&\frac{v(G)}{2\ell}\left(\min_{x\in V(G)} \deg(H_x)\right)\left(\min_{x\in V(G)} \text{number of } (2\ell-2m-1)\text{-paths in } H_x\right)\\
&\ge \frac{n}{2\ell}\left(\frac{c_1^{m+1}}{10}n^{1/m}\right)\left(\frac{9c_1^{(2\ell-2m)(m+1)}}{10^{2\ell-2m}c_2}n^{(2\ell-m-1)/m}\right)\\
&= \frac{9}{2\ell c_2}\left(\frac{c_1^{m+1}}{10}\right)^{2\ell-2m+1} n^{2\ell/m}.
\end{aligned}$$

□

4 Proof of main theorem

To count the number of longer cycles, we observe that H_x has all the nice properties we wanted, namely bipartite, almost regular and large girth, we can apply Proposition 8 to H_x and get many short cycles in H_x. From them, we obtain a lot of paths in H_x, and each of them corresponds to many longer cycles in G as in Section 3. These longer cycles give many longer paths in H_x and again, each of these paths corresponds to many even longer cycles in G_x. Eventually, we have Theorem 2.

For simplicity, we will assume that m is even from now on. For odd m, one can proceed similarly.

Changing the parameters in Proposition 8, the number of 2ℓ-cycles in H_x is at least

$$\frac{9}{2\ell c_2}\left(\frac{c_1^{(m+1)^2}}{10^{m+2}}\right)^{2\ell-2m+1} n^{2\ell/m},$$

for $2\ell \in L_0 := \{3m, 3m+2, 3m+4, \ldots, 4m\}$, and so the number of paths of length $2\ell - m \in \{2m, 2m+2, 2m+4, \ldots, 3m\}$ in H_x is at least

$$\frac{9}{c_2}\left(\frac{c_1^{(m+1)^2}}{10^{m+2}}\right)^{2\ell-2m+1} n^{2\ell/m}.$$

Then, for $2\ell \in L_0$, the number of $((2\ell-m)+2m+2)$-cycles in G is at least

$$\frac{n}{2\ell+m+2}\frac{c_1^{2(m+1)}}{100}n^{2/m}\frac{9}{c_2}\left(\frac{c_1^{(m+1)^2}}{10^{m+2}}\right)^{2\ell-2m+1} n^{2\ell/m}$$
$$= \frac{9}{(2\ell+m+2)c_2}\frac{c_1^{2(m+1)}}{100}\left(\frac{c_1^{(m+1)^2}}{10^{m+2}}\right)^{2\ell-2m+1} n^{(2\ell+m+2)/m},$$

or for $2\ell \in L_1 := \{4m+2, 4m+4, \ldots, 5m+2\}$, the number of 2ℓ-cycles in G is at least $\alpha_\ell n^{2\ell/m}$, where α_ℓ is a positive constant depending on m, c_1, c_2, ℓ only. Repeating the same argument with the sets $L_j := \{3m+j(m+2), 3m+j(m+2)+2, \ldots, 4m+j(m+2)\}$, the proof of Theorem 2 is completed.

References

[1] C. T. Benson, *Minimal regular graphs of girths eight and twelve*, Canad. J. Math. **18** 1966 1091-1094.

[2] J. A. Bondy and M. Simonovits, *Cycles of even length in graphs*, J. Combin. Theory Ser. B **16** (1974), 97-105.

[3] W. G. Brown, *On graphs that do not contain a Thomsen graph*, Canad. Math. Bull. **9** 1966 281-285.

[4] P. Erdős, *Extremal problems in graph theory In Proc. Symp. Theory of Graphs and its Applications*, page 2936, 1963.

[5] I. Reiman, *Über ein Problem von K. Zarankiewicz*, Acta. Math. Acad. Sci. Hungar. **9** 1958 269-273.

[6] B. Sudakoov and J. Verstraëte, *Cycle lengths in sparse graphs*, Combinatorica 28 (2008), no. 3, 357372.

[7] R. Wenger, *Extremal graphs with no C4's, C6's, or C10's*, J. Combin. Theory Ser. B 52 (1991), no. 1, 113-116.